



### Summer News

**The Westpoint Southern region would like to thank you all for working with us during another extremely busy year!**

#### Veterinary students

To support and encourage future farm vets, we are a partnership practice for the Surrey Veterinary School. Final year students from the vet school will be completing their month-long farm rotations with us. They will be on farm with our vets learning day 1 skills as well as an understanding of current farming systems. Please help us support them and don't worry, additional time on farm for teaching will not be charged.

#### **Ashford**

After a busy move we are now settled in our new practice premises and our pleased to be sharing it with **FarmSupplies**. Feel free to drop in and see us at Unit 11, St. Johns Court, Sevington Business Park, Ashford. We are open for enquiries and medicine pick-ups Monday to Friday 8am -5pm.

The farm vet team at Ashford consists of Kathy, Alex, Jack, Sharminda and Alice, sadly Jess has moved to pastures new as a farm animal vet in Devon to be closer to home. All our vets are dedicated farm vets and as well as the routine work, they all have areas of special interest. Jack is currently studying for a Diploma in Bovine Reproduction, a two-year master's program, to further his knowledge in this area and Sharminda is carrying out a certificate in advanced veterinary practice in sheep. The skills they learn will be of great benefit to the practice and the clients.

#### BVD Stamp It Out!

Get involved! We are really pleased that 25 farms have already joined the BVD project for breeding cattle herds. It is a funded project to help farmers to take action in BVD control, we would like to help more farms to join and take action. So please contact us for more information if you are interested in joining this nationwide project.

#### Barren ewes post lambing?

Toxoplasmosis and Enzootic abortion are common causes of abortion in sheep and have significant effects on productivity. If the status of your flock is unknown, we can blood test 6-8 barren ewes post lambing and screen them for these diseases. For flocks of over 100 breeding ewes we can use the Flockcheck test, sponsored by MSD. For more information please contact the Ashford practice.

#### **Sevenoaks and Heathfield**

We are very excited that in November last year vet Maria Sara Santos joined our practice. Her areas of interested are internal medicine and she is definitely a great addition to the team. Currently, the vets are Rui d'Orey Branco, Practice Principal, Sarah O'Reilly, Rebecca Rehage, and Maria Sara Santos. Our receptionist is Jana Saunders. The Heathfield office is currently open sporadically as Sandra Butchart has recently retired. The office will be open as usual as soon we have appointed a new receptionist.

#### Flock Health Club

We are pleased to announce the start of our Flock Health Club.

For a small monthly fee, a range of benefits will be available at a discounted rate; including a full Red Tractor compliant health plan, and a social event per year. For more information please contact Sarah on sarah.oreilly@westpointfarmvets.co.uk.

#### Small Holder Club

We are very excited to announce that we had the first meeting for the Smallholder Club and we believe everyone enjoyed it! This club will suit farmers/owners that gives names to their animals. We have 3 to 4 meetings a year and an allocated vet with a special interest in the subject. For more information contact Maria Sara on mariasara.santos@westpointfarmvets.co.uk.

#### City Farms Club

We have several public farms with a huge variety of species. The main goal of this club is to help the farmers to develop competencies in some of the emergency situations and increase the networking across these farms. For more information contact Rebecca on rebecca.rehage@westpointfarmvets.co.uk.

#### Alpacas Club

The alpaca population is growing and these cute animals need special treatment. We are creating a club to increase the sharing of knowledge and provide a more detailed service to meet the needs of these exotic animals. There will be a set of talks and services specific to alpacas. For more information contact Rebecca on rebecca.rehage@westpointfarmvets.co.uk.

#### Pro Dairy Training

We are currently running a set of 10 courses throughout a 12 month period. The objective is to train farmers to become confident in different skill sets needed on farm. These are very practical courses with topics such as Artificial Insemination, Managing difficult calving, Milk parlour routine, Nutrition, Medicine usage and administration. For more information contact Rui on rui.branco@westpointfarmvets.co.uk.

#### Sevenoaks Upcoming Events

19th June, time TBC – Flock Health Club: An overview of Infectious diseases (the iceberg diseases).  
24th June, time TBC – Pump Clinic and Oral Fluids Therapy  
10th of July, time TBC – Sheep Mineral and Trace Elements – Bimeda

#### **Summer Show Season**

Through the show season we will be at the following shows providing vet care and we will have a stand at some of them. Please come find us for a chat and a cup of tea.

- Oxted and Edenbridge Show – 25th-26th August
- Findon Sheep Fair – 14th September
- West Grinstead Ploughing Match – Saturday 21st September
- East Kent Ploughing Match – Wednesday 25th September

Hopefully we will see you at one of the approaching shows, but if not, we are always on the end of the phone for advice, queries, or to take any feedback that you may have for us.

***Have a great summer!***

# 5 Steps to a Successful Unpacking Season

by Ami Sawran BVSc MRCVS

Unpacking, though exciting, can be a little tiring, and seems to come around quickly each year, so in this article, I hope to relay some gentle reminders and helpful tips to ensure that things run as smoothly as possible for all alpaca and llama owners gearing up for the season.

## 1. Getting Kitted Out

Something that can be genuinely life-saving is having a birthing kit prepared, to be grabbed at a moment's notice. In that kit, you may have medicines ranging from vitamins, to pain relief/anti-inflammatory, to antibiotic. Your vet may not be able to dispense these medications 'as and when' without a visit (as they must certify your animals 'under their care'), so it pays to ensure you have had a consultation with your vet, so you may keep medicines on the premises for when you need them most.

As for what else should feature in the kit, consider small lambing ropes, a lambing snare (protected wire to help safely align the head), disinfectant, clean water, scissors, towels, a clamp for a bleeding umbilicus, an oesophageal feeding tube (such as for lambs), long-armed gloves, iodine for naval dipping, lots of lubricant (lube bottles with a long spout are most useful), a headtorch, your vet's telephone number and some plastic sheeting (in case of a uterine prolapse, you can keep it clean on the sheet until a vet arrives). If you are unsure how to safely use ropes and a snare, ask for a demonstration from you vet beforehand.



## 2. Consider Plasma Banking

Though we hope that all cria will be born fighting fit, and that the hembra will have enough good quality colostrum for them, it sometimes transpires that the cria does not get enough colostrum, or the colostrum is not of sufficient quality to convey vital antibodies to diseases. In this case, a cria may have failure of passive transfer (FPT) of immunity. FPT cria are best treated with an intravenous infusion of plasma that has been spun down from the blood of vaccinated, healthy, usually male alpacas on your farm. Plasma is also indicated for premature cria after a difficult birth, for those with congenital defects such as cleft palate, and in cases of mismothering. Ask your vet if they can provide this service prior to unpacking, to allow you to stockpile an adequate supply of frozen plasma, which can be stored for up to 2 years.

## 3. The Back-up Plan

If plasma banking is not available to you, there are still ways you can supplement cria if they are not feeding appropriately. Cria must consume 10-20% of their bodyweight in the first 24 hours of life (over two hourly feeds), with a golden window for absorption of vital antibodies in the first 8 hours. Keep track of the cria's weight from the day of birth onwards. If it appears the cria is not getting this from the hembra (sometimes indicated by vocalising, increased attempts to feed, or not swallowing during nursing), then you may need to bottle, or even tube feed an alternative. Goats colostrum and milk is the most common alternative, though if you are sourcing this from a farm, please ensure that it is from a Johne's disease free

source. Again, if you are unsure how to safely tube feed, as for a demonstration ahead of time.

## 4. When To Call A Vet

Though many camelid owners are very experienced with unpacking, it's worth knowing when veterinary intervention is required. As we expect most unpacking to occur between 10am and 2pm, anything outside that time is worth investigating. Keep in mind once the hembra is actively labouring, some progress should be made every 15 minutes; and cria should be delivered within 45 minutes. If, for example, only one leg is visible for more than 15 minutes, this warrants gentle investigation with a well-lubricated, gloved hand. Check whether the other leg is simply tucked up against the pelvis and holding up the process – can this be carefully brought round, guarding the delicate walls of the uterus from the foot? Is the head within reach? Labour cannot progress without the head being in position; upright and between the feet. Shoulder lock is the most common cause of cria becoming stuck, which can require gentle rotation of the cria. If you do not feel able to correct the cria's position, this may need to be done under epidural by your vet. If you are unsure of whether you have a back or front foot, remember that the first two joints of the foot bend the same way on the forelimbs (making a 'U' shape), and opposite ways on hindlimbs (making a 'Z' shape). If you have a cria coming backwards (back legs, or rump first), it may be sensible to speak with your vet, as backwards delivery is more difficult and can compromise the cria's breathing. If you are sure that labour is occurring but the hembra isn't open normally, she may have a uterine torsion, which requires veterinary intervention. If you feel that everything is now in the correct position, back away and allow the hembra to progress naturally – pulling on a cria can cause damage to the uterus and vaginal tissues, or bring about a prolapse. If ever in doubt, speak with your vet, who will be able to guide you – the camelid reproductive tract is fragile, so it is better to err on the side of caution always.

## 5. The Aftermath

After the cria is delivered, check membranes and mucus are cleared from the nose. Do not be tempted to 'swing' the cria or hang it over a gate – this compresses the lungs. Place the cria so it is lying on its sternum (breast bone) and rub the sides of the chest with a towel to stimulate breathing. Cleanly and carefully check for any damage (tears or bruising) to the vaginal tissue – this may require veterinary examination, or the administration of antibiotics and pain relief. If you do note damage and excess bleeding, applying pressure with clean, damp gauze can help before your vet arrives. Check also for a rare twin. Don't be tempted to pull on the placenta just yet – this should be passed in its entirety within 6 hours of birth. If this has not happened, or you have checked it and it is not entire (it is essentially a sack that envelopes the whole cria, so shouldn't have jagged pieces missing), then you may need your vet to check the hembra and potentially administer medicines to help her pass the remnants.

**Preparation, as always, is key to the smooth running of the unpacking season, but in the event that you are unsure about how to handle unpacking, from labour to lifelong care, don't hesitate to contact your veterinarian to put a bespoke birthing and health plan in place.**

**Westpoint provide a plasma generation service. If you would like to discuss this further, or find out if your location is eligible, please call your local Westpoint practice.**



# What does IBR look like on my farm?

by Tony O'Loughlin BVSc MRCVS

**Infectious Bovine Rhinotracheitis – IBR – is a herpes virus that can cause respiratory disease in cattle of all ages or reproductive disease in breeding females. It is caused by Bovine Herpes Virus 1 – BHV 1 – and has a worldwide distribution. Initially the virus was recognised as causing severe upper respiratory tract disease in cattle – classically these would have presented with a severe mucopurulent discharge from the nose and eyes – lots of thick pus! The animals had a very high temperature and they would have shown coughing, increased breathing and generally off colour and inappetent. The disease would have spread rapidly on a farm and mortality could be very high.**

Today we tend to see a more subtle disease pattern. There can be a background cough with slight nasal and ocular discharge. We then start to see the more subtle signs of an effect on production – lower milk yield and reduced fertility. These production effects can be missed and represent a “drag” on farm performance.

IBR is spread by carrier animals. Any animal that becomes infected will become a latent carrier – as humans do with the herpes simplex virus that causes cold sores. These latently infected animals have no clinical signs but can start to re-excrete at any time following a period of stress or an immune challenge. These latently infected animals are the source of infection in the herd. For an IBR free herd these animals are the greatest danger.

IBR can be controlled and avoided. The first step is to establish your IBR status with your vet – this can be done via blood samples or a bulk tank sample. We look for antibodies proving exposure to IBR virus. If antibodies are present, and there are no vaccinated animals present, then the animals have been exposed to IBR and will have latent carriers present.



## IBR Negative

If you are fortunate enough to be IBR negative, then you must take steps to protect that status – negative also means the herd is susceptible to infection. **Biosecurity is key here.** As discussed above the biggest risk to your herd is from purchasing latent carriers – unless you sample incoming animals you won't know their health status. You should do a biosecurity risk assessment with your vet as part of your herd health control. This will include assessing risks from neighbouring cattle, farm visitors etc. You may choose vaccination as one control option if your risk assessment of infection is high.

Or you can choose to buy from an IBR Accredited herd. There are CHeCS accreditation schemes for the common infectious diseases of cattle. Cattle herds can be accredited free from IBR, BVD, Leptospirosis, Johnes, Neospora and TB. The schemes for each disease run separately so you can choose what suits you.

## IBR Positive

If you are IBR positive, then your aim is to limit the effects of the infection in the herd – reduce the clinical signs – and to prevent the development of latent carriers. You can choose to sample all the individuals in the herd and to cull the positive animals. This can be expensive and may not be economically viable.

**Vaccination** will limit disease transmission between animals, helping to protect uninfected animals. It will also prevent the development of latency thus slowing spread within a herd. Carried out correctly, vaccination should also reduce the intensity and duration of clinical signs.

Using a live marker vaccine means surveillance testing can continue to be carried out after vaccination, as vaccinated animals can be distinguished apart from animals exposed to wild type IBR.

## Summary

Every cattle farm – dairy or beef – should know their IBR status as the effects of the disease can be subtle and production limiting. The principles of infectious disease control then apply...

- Know your health status for the important diseases
- Assess your biosecurity to keep disease out and to limit disease spread
- Implement prevention and/or control measures
- Work with your vet.

**Contact your local Westpoint practice for more information on IBR control and eradication in your dairy or beef herd and the benefits of developing a bespoke herd health plan.**



## Fly Control Offers

farmacy 



PRODUCT	WAS	NOW
Butox Swish 12L	£360.40	<b>£355.00</b>
Crovect 2.2L	£47.10	<b>£43.00</b>
Dectospot 2.5L	£175.50	<b>£151.00</b>
Deltanil 2.5L	£163.00	<b>£155.00</b>
Ectofly 5L	£64.56	<b>£59.00</b>
Electron Fly Tags 20 Pack	£53.20	<b>£49.00</b>
Flypor 12L + Applicator	£305.29	<b>£278.00</b>
Spotinor 2.5L	£194.72	<b>£152.50</b>

now you can collect farmacy  
**REWARD POINTS**  
everytime you order!

Quote code **WNL**  
[www.farmacy.co.uk/WNL](http://www.farmacy.co.uk/WNL)

Tel: **01306 628215**

Use Medicines Responsibly ([www.noah.co.uk/responsible](http://www.noah.co.uk/responsible)) Always seek advice on the correct use of this or alternative medicines from the medicine prescriber. PLEASE NOTE: A prescription issued by a UK registered vet (MRCVS) is required before some items can be dispatched. Cannot be used in conjunction with any other offers or discounts. Offer subject to availability. Offer ends 30th June 2019.

# Tackling Lameness in Sheep with the Five-Point Plan

by Emily Francis BVM&s MRCVS BSc

Tackling lameness in sheep has become a real priority in recent times and much research has been undertaken to determine the most effective ways to control and prevent it. The Farm Animal Welfare Council have set a target to reduce the number of sheep treated, per flock per year, to 2% by 2021. There is also a huge emphasis on reducing antibiotic use on farm; and as lameness treatment is a significant contributor, farmers are being encouraged to tackle their lameness issues head on.

The five-point plan was developed by combining previous research with real on farm experience. The five points work together and have 3 ultimate goals: build resilience, establish immunity and reduce the disease challenge.

## Where to Start?

Firstly, the best way to assess whether any changes you undertake are making a difference, is to know what your starting lameness prevalence is. The number of animals treated over a particular time period is an easy one to monitor but requires strict recording of all the animals treated day to day. Secondly, it is vitally important to correctly identify the cause of the lameness on your farm, as treatment and prevention methods may vary. Footrot causes approximately 80% of lameness and is therefore the focus of the five-point plan. However, it may be that contagious ovine digital dermatitis (CODD), a combination of both footrot and CODD or other non-infectious causes may be causing the issues. Some tips on correct identification include:

- Footrot has a characteristic foul smell. Lesions start as a reddening in between the digits (scald) before an underrunning of the horn is seen. CODD does not produce this characteristic smell, and the hoof wall tends to be lost from the outer coronary band.
- Use published guides from organisations such as AHDB Beef and Lamb Better Returns to make more accurate decisions.

The points of the plan should be discussed with your vet to aid the implementation of changes that best suit your farm and type of lameness.

## Putting the Five-Point Plan Into Action

- 1. Culling repeatedly affected animals** works to prevent chronically lame individuals from spreading the bacteria and acting as reservoirs of infection to other animals. Sheep that have been treated for footrot more than twice in one season should be culled before the next. However, this requires significant commitment, as she may be a particularly good breeding ewe. Some things to consider when culling:
  - Culling hard in the first two years of the five-point plan should result in a lower lameness prevalence over the following years so the culling rate for lameness should decrease again with time.
  - If there is no current culling policy for lame animals then using a higher cut off e.g. three treatments in a season may still help.
  - This approach relies on good recording and so tagging repeat offenders and marking the leg they are lame on at each treatment should be undertaken.

- 2. The quarantine of incoming animals** is particularly important, as purchased stock act as a source of many diseases including

CODD and new strains of footrot. Ideally all animals are bought in from farms with low lameness levels and quarantined for at least 4 weeks. In this time, all cases of lameness are treated and new cases monitored so that all animals are sound before mixing with the main flock. One option is to run new animals through a footbath before turning them onto your isolation pasture.



- 3. Treating clinical cases early** has been shown to dramatically reduce lameness levels. This is because there is not only a greater chance of curing an animal but because it also reduces the time that lame animal acts as a reservoir of infection.

- Footrot is caused by the bacteria so antibiotics are appropriate.
- Lame animals should be treated within 3 days of identification.
- Animals with early footrot (scald) should have all four feet sprayed with oxytetracycline spray (blue) more severe cases with underrun horn should be treated with a long acting injectable oxytetracycline. Most long acting preparations only last 2-3 days and so re-treatment after the appropriate time is fundamental to successful treatment.
- Ensure animals are dosed for the correct weight.
- **Preventative and therapeutic foot trimming should not be undertaken** as it has been shown to delay recovery.
- Isolate repeat cases to prevent spread to young lambs in particular. The use of "lame pen/field" closer to the farm aids treatment and monitoring of lame sheep.

- 4. Avoid the spread of infection** by thinking about when and how you handle the flock:

- Footbathing is a very useful tool in treating a scald outbreak in lambs and for disinfecting the feet of incoming animals. It does not treat footrot so all clinically lame animals should be separated and not run through the bath with the others.
- Can your handling system be moved to clean underfoot areas each time? Is it thoroughly cleaned and disinfected between uses?
- Lime can be used in holding pens or around troughs to reduce bacterial load in areas of high traffic.

- 5. Vaccinating** animals not only raises an individual's immunity to footrot but results in a whole flock immunity due to the dramatic decrease in the number of animals spreading infection. The commercially available vaccine (Footvax: MSD Animal Health) is a subcutaneous injection, given as a two-dose primary course and then boosted before perceived risk periods occur. Its use should be discussed with your vet to determine the best vaccination programme for your farm and optimum results from the vaccine are seen when other points in the plan are used in conjunction.

**The flexibility of this plan means that not all of the points need to be worked on at once, but used to identify those that are easier short-term goals or that need immediate attention. Commitment is needed to reduce lameness using the five-point plan however, with persistence, this approach can result in significant improvements in your flock.**

## Westpoint Farm Vets

**Sevenoaks** Park Farm Stables, Park Farm, Brasted, Nr Sevenoaks, Kent, TN16 1LL | 01959 564383  
**Heathfield** Monkhurst Offices, Sandy Cross Lane, Heathfield, East Sussex, TN21 8QR | 01435 869047  
**Ashford** Unit 11 St Johns Court, Foster Road, Ashford, Kent, TN24 0SJ | 01306 628208